



Contents lists available at ScienceDirect

Seizure: European Journal of Epilepsy

journal homepage: www.elsevier.com/locate/seizure

Erratum



Erratum to “Characteristics of post-ictal headaches in patients with epilepsy: A longitudinal study” [Seizure: Eur J Epilepsy 81 (2020) 244–249]

Ana Letícia F. Caprara ^{a,*}, Jamir P. Rissardo ^a, Martim T.B. Leite ^a, Juliana O.F. Silveira ^a, Paulo Gilberto M. Jauris ^a, Josi Arend ^b, Aline Kegler ^c, Luiz Fernando F. Royes ^{d,e}, Michele R. Figuera ^a

^a Health Sciences Center, Department of Neuropsychiatry, Federal University of Santa Maria, RS, Brazil

^b Postgraduate Program in Pharmacology, Federal University of Santa Maria, RS, Brazil

^c Center for Natural and Exact Sciences, Postgraduate Program in Biological Sciences, Federal University of Santa Maria, RS, Brazil

^d Toxicological Biochemistry, Federal University of Santa Maria, RS, Brazil

^e Physical Education and Sports Center, Exercise Biochemistry Laboratory (BIOEX), Federal University of Santa Maria, RS, Brazil

The publisher regrets that the following errors are present in the published article:

1) The names of the second and tenth authors are incorrect; the correct names are given above.

	Migrainous N = 45 (Group 1)	Tension-type N = 77 (Group 2)	Mixed N = 18 (Group 3)	Odds Ratio ¶	95 % CI	p-value
Seizure onset type						
Generalized onset	25 (55.5)	46 (59.7)	14 (77.8)	1.28	0.97 – 1.34	0.421
Focal onset	19 (42.2)	29 (37.7)	4 (22.2)	1.13	0.74 – 1.01	0.633
Focal to bilateral tonic-clonic	14 (31.1)	16 (20.8)	1 (5.5)	0.91	1.15 – 1.29	0.707
Unknown	1 (2.2)	2 (2.6)	0	–	–	–
Etiology						
Structural	12 (26.7)	23 (29.9)	7 (38.9)	2.15	1.71 – 1.92	0.543
Unknown	20 (44.4)	35 (45.4)	6 (33.3)	1.55	0.83 – 1.69	0.868
Infectious	11 (24.4)	17 (22.1)	4 (22.2)	1.21	0.54 – 1.23	0.997
Genetic	2 (4.4)	2 (2.6)	1 (5.5)	1.12	1.56 – 2.33	0.554
DRE (n = 33)	21 (46.7)	9 (11.7)	3 (16.7)	1.45	1.33 – 1.61	0.004*
Non-DRE (n = 107)	24 (53.3)	68 (88.3)	15 (83.3)	1.73	1.45 – 1.82	0.003*
AED polytherapy (n = 58)	22 (48.9)	30 (38.9)	6 (33.3)	1.84	0.91 – 2.65	0.423
AED monotherapy (n = 82)	23 (51.1)	47 (61)	12 (66.7)	0.984	0.652 – 1.04	0.562
Valproate	7 (15.5)	15 (19.5)	3 (16.7)	–	–	–
Topiramate	3 (6.7)	9 (11.7)	1 (5.5)	–	–	–
Carbamazepine	6 (13.3)	11 (14.3)	5 (27.8)	–	–	–
Lamotrigine	4 (8.9)	8 (10.4)	2 (11.1)	–	–	–
Other	3 (6.7)	4 (5.2)	1 (5.5)	–	–	–
Valproate	7 (15.5)	15 (19.5)	3 (16.7)	–	–	–
Migraine with aura	2 (4.4)	5 (6.5)	0	1.15	1.23 – 1.36	0.622
Migraine without aura	15 (33.3)	42 (54.5)	7 (38.9)	1.26	1.05 – 2.87	0.023*
Probable migraine	5 (11.1)	3 (3.9)	2 (11.1)	0.85	0.71 – 3.24	0.570

DOI of original article: <https://doi.org/10.1016/j.seizure.2020.08.001>.

* Corresponding author at: Av. Roraima n° 1000, Cidade Universitaria, Bairro Camobi, Santa Maria, RS, Brazil.

E-mail address: ana.leticia.fornari@gmail.com (A.L. F. Caprara).

<https://doi.org/10.1016/j.seizure.2020.09.005>

Available online 18 September 2020

1059-1311/© 2020 British Epilepsy Association.

Published by Elsevier Ltd.

This article is made available under the Elsevier license

(<http://www.elsevier.com/open-access/userlicense/1.0/>).

Group	Variable	Dependent variable			
		Seizures in the last follow-up			
		Odds ratio	95 % CI	p value	Parameter estimate ^a
Post-ictal headache (+)	Family history of migraine	1.696	1.372 to 2.096	0.032*	0.015
	Drug-resistant epilepsy	1.169	1.135 to 2.146	0.029*	0.002
	Months since last visit	1.464	1.243 to 2.888	0.035*	0.006
	Seizure Onset Type “Generalized”	1.527	1.114 to 1.668	0.004*	0.001
	Family history of seizures	1.001	0.851 to 1.530		
	School performance (poor)	1.021	0.989 to 1.244		
	Migraine	1.539	0.867 to 1.622		
	AED monotherapy	1.015	0.763 to 2.691		
	AED polytherapy	1.058	0.952 to 2.029		
	Treatment adjustment	1.013	0.875 to 1.384		
Post-ictal headache (-)	Frequency of seizures before treatment	1.005	1.225 to 3.641		
	Family history of seizures	0.884	0.547 to 1.973		
	Drug-resistant epilepsy	0.985	0.622 to 0.991	0.032*	-0.025
	Months since last visit	0.632	0.854 to 1.751		
	Seizure Onset Type “Generalized”	0.739	0.421 to 0.895	0.009*	-0.013
	Family history of migraine	0.778	0.545 to 1.215		
	Generalized (Focal as baseline)	0.692	0.531 to 1.790		
	AED monotherapy	0.815	0.967 to 1.741		
	AED polytherapy	0.744	0.854 to 1.298		
	Treatment adjustment	0.732	0.525 to 1.642		
	Frequency of seizures before treatment	0.821	0.774 to 2.658		
	Migraine	0.992	0.855 to 1.249		

2) The alignment of the data in Tables 2 and 3 is incorrect. The corrected Tables are provided below.

Table 2 Clinical characteristics of subjects according with type of PIH.

Abbreviations: N: Number; () represent percentages, CI: Confidence Interval; *: P < .05.

¶: For the bivariate analysis, only Groups 1 and 2 were compared.

Table 3 Estimated Fixed Effects of Predictors in the Multilevel Linear Regression Model.

^aParameter estimates represents the effects of a determined predictor on the population.

Variables that were not included in the model have empty entries.

Abbreviations: CI: Confidence Interval; *: P < .05.

The publisher would like to apologise for any inconvenience caused.